



Annual distribution of black cardinal fish lineage

FD0240_1; FD0241_1; FD0242_2; FD243_2; FD244_2

1. Electronic databases were used to generate initial maps of species distribution.
 - a. Commercial fishing returns (larger vessels): **TCEPR** database. All records from 1 October 1989 to 30 September 2005 were extracted on 17 October 2005. Data were used to estimate mean annual catch and catch rate (kilograms per kilometre towed) in 0.25 degree rectangles. Only the top five species caught are reported on these forms so information on the absence of a species is not available.
 - b. Commercial fishing returns (smaller vessels): **CELR** database. All records from 1 October 1989 to 30 June 2003 were extracted on 15–17 July 2003. Data were used to estimate mean annual catch in statistical areas. Information from statistical areas 1–10 was down-weighted because of likely mis-recording of Fishstock instead of statistical area. Only the top five species caught are reported on these forms so information on the absence of a species is not available.
 - c. Scientific observer records from larger vessels: **obs** database. All records from 1 March 1990 to 30 September 2005 and stored in the new data format were extracted on 20 October 2005. Data were used to estimate mean annual catch and catch rate (kilograms per kilometre towed), and proportion of tows that caught the species, in 0.25 degree rectangles.
 - d. Research bottom trawl records: **fish_comm** database. This database is a groomed version of the research trawl database **trawl**. All records from 2 September 1978 to 30 September 2005 were extracted on 19 May 2006. Data were used to estimate total catch, proportion of tows that caught the species, and catch rate (kilograms per kilometre towed) in 0.25 degree rectangles.
 - e. Russian research bottom trawl records: **trawl** database. These data are a subset of the research trawl database **trawl**. All records were extracted on 9 August 2003. Data were used to determine the presence of this species north of 37 °S. Because the data are old (the most recent record was 1987), and there are problems with species identifications, these data were given low weighting.
 - f. Museum of New Zealand Te Papa records of this species based on voucher specimens held in their collection were searched for distributional information that added to the distributional ranges determined from other databases.
 - g. Databases of commercial tuna longline catches (**TLCER**), observer records from tuna longlines (**I_line**), recreational fishing (**rec_data**), and aerial sightings (**aer_sight**) were not used as they contained

no records of this species, or the number of records was too small to provide useful additional distributional information.

2. Literature sources were searched for distributional information that added to the distributional ranges determined from databases.
 - a. Unpublished electronic bibliography of New Zealand fishes compiled by L. J. Paul and held on a NIWA computer.
 - b. Aquatic Sciences and Fisheries Abstracts.
 - c. *New Zealand Professional Fisherman* and *Seafood New Zealand* for 1986–2002.
 - d. *New Zealand Fishing News* for 1998–2002.
 - e. Scientific papers, unpublished reports and university theses available to the expert who prepared the distributional layers.
3. Other sources.
 - a. Nil.
4. Summary
 - a. Maps generated from the electronic databases were provided to an expert scientist who integrated this information with other information from the literature, and expert opinion, and produced hand-drawn distributional zones on a template map containing depth contours at 250 m, 500 m, and 1000 m. These maps were then digitised and imported into a GIS software package as layers. The areas of the zones were calculated, and the layers were linked to attribute and metadata files.
 - b. The primary sources of distributional data for black cardinal fish were TCEPR, CELR, and fish_comm databases.
 - c. Black cardinal fish is widely distributed in New Zealand waters. Although several species of cardinal fish occur, black cardinal fish is the only species that reaches a marketable size, and is therefore the only commercial species at present. It is often associated with seamount features, and extends out from the continental shelf and slope along ridges north of North Island. The extent of northern and southern distributions is uncertain, and it appears to be absent from the eastern regions of the Southern Plateau. It covers a wide depth range of 250–1500 m, but is most common between 500 m and 1000 m.

5. References

The following sources provided useful information on the distribution of this species. This is not an exhaustive list of all references to the species.

Anderson, O.F.; Bagley, N.W.; Hurst, R.J.; Francis, M.P.; Clark, M.R.; McMillan, P.J. (1998). Atlas of New Zealand fish and squid distributions from research bottom trawls. *NIWA Technical Report 42*. 303 p.

Clark, M.R.; King, K.J. (1989). Deepwater fish resources off the North Island, New Zealand: results of a trawl survey, May 1985 to June 1986. *New Zealand Fisheries Technical Report No. 11*. 56 p.

- Dunn, M.R. (2005). Descriptive analysis of catch and effort data from New Zealand black cardinalfish (*Epigonus telescopus*) fisheries for the fishing years 1979–80 to 2002–03. *New Zealand Fisheries Assessment Report 2005/32*. 48 p.
- Field, K.D.; Tracey, D.M.; Clark, M.R. (1997). A summary of information on, and an assessment of the fishery for, black cardinalfish, *Epigonus telescopus* (Risso, 1810) (Percoidei: Apogonidae). New Zealand Fisheries Assessment Research Document 97/22 (unpublished report held in NIWA library, Wellington).
- Field, K.D.; Clark, M.R. (2001). Catch-per-unit-effort (CPUE) analysis and stock assessment for black cardinalfish (*Epigonus telescopus*) in QMA 2. *New Zealand Fisheries Assessment Report 2001/23*. 22 p.
- Francis, M.P.; Hurst, R.J.; McArdle, B.H.; Bagley, N.W.; Anderson, O.F. (2002). New Zealand demersal fish assemblages. *Environmental Biology of Fishes* 65: 215-234.
- Phillips, N.L. (2002). Descriptive and catch-per-unit-effort (CPUE) analyses for black cardinalfish (*Epigonus telescopus*) in QMA 1. *New Zealand Fisheries Assessment Report 2002/55*. 54 p.