



Lineage

Summer distribution of kahawai

1. Electronic databases were used to generate initial maps of species distribution.
 - a. Commercial fishing returns (larger vessels): **TCEPR** database. All records from 1 October 1989 to 30 June 2003 were extracted on 16 July 2003. Data were used to estimate mean annual catch and catch rate (kilograms per kilometre towed) in 0.25 degree rectangles. Only the top five species caught are reported on these forms so information on the absence of a species is not available. Records of kahawai from depths greater than 150 m, and on the Snares Shelf and Campbell Plateau, are probable mis-identifications or mis-codings of other species, and were ignored (kahawai has never been recorded from these regions in research trawl tows).
 - b. Commercial fishing returns (smaller vessels): **CELR** database. All records from 1 October 1989 to 30 June 2003 were extracted on 15–17 July 2003. Data were used to estimate mean annual catch in statistical areas. Information from statistical areas 1–10 was down-weighted because of likely mis-recording of Fishstock instead of statistical area. Only the top five species caught are reported on these forms so information on the absence of a species is not available. Records of kahawai from depths greater than 150 m, and on the Snares Shelf, Campbell Plateau and Bounty Plateau, are probable mis-identifications or mis-codings of other species, and were ignored (kahawai has never been recorded from these regions in research trawl tows).
 - c. Scientific observer records from larger vessels: **obs** database. All records from 1 March 1990 to 30 June 2003 and stored in the new data format were extracted on 28 July 2003. Data were used to estimate mean annual catch and catch rate (kilograms per kilometre towed), and proportion of tows that caught the species, in 0.25 degree rectangles.
 - d. Research bottom trawl records: **fish_comm** database. This database is a groomed version of the research trawl database **trawl**. All records from 2 September 1978 to 3 September 1997 were extracted on 15–16 July 2003. Further surveys have been added to **trawl** and **fish_comm** since 1997, but were not used because they have not been properly groomed for species identification and positional errors. Data were used to estimate total catch, proportion of tows that caught the species, and catch rate (kilograms per kilometre towed) in 0.25 degree rectangles.
 - e. Recreational fishing database: **rec_data**. All records were extracted on 24 July 2003. Data were used to determine the presence of a species in a variety of statistical reporting areas.

- f. Aerial sightings database: **aer_sight**. On 5 August 2003, data were extracted for 1976 onwards (for 0.5 degree squares) and for 1 January 1986 onwards (for actual positions). Data were used to estimate total tonnage, number of schools, and tonnes per hour of flying.
 - g. Museum of New Zealand Te Papa records of this species based on voucher specimens held in their collection were searched for distributional information that added to the distributional ranges determined from other databases.
 - h. Databases of commercial tuna longline catches (**TLCER**), and observer records from tuna longlines (**l_line**) were not used as they contained no records of this species, or the number of records was too small to provide useful additional distributional information. Records from Russian trawl surveys (**trawl**) were not used because they were historic (pre 1987), and species identification (i.e., separation of golden and horse mackerels) is considered unreliable.
2. Literature sources were searched for distributional information that added to the distributional ranges determined from databases.
 - a. Unpublished electronic bibliography of New Zealand fishes compiled by L. J. Paul and held on a NIWA computer.
 - b. Aquatic Sciences and Fisheries Abstracts.
 - c. *New Zealand Professional Fisherman* and *Seafood New Zealand* for 1986–2002.
 - d. *New Zealand Fishing News* for 1998–2002.
 - e. Scientific papers, unpublished reports and university theses available to the expert who prepared the distributional layers.
 3. Other sources.
 - a. Nil.
 4. Summary
 - a. Maps generated from the electronic databases were provided to an expert scientist who integrated this information with other information from the literature, and expert opinion, and produced hand-drawn distributional zones on a template map containing depth contours at 250 m, 500 m, and 1000 m. These maps were then digitised and imported into a GIS software package as layers. The areas of the zones were calculated, and the layers were linked to attribute and metadata files.
 - b. The primary sources of distributional data for kahawai were the **aer_sight**, **TCEPR**, **CELR**, and **fish_comm** databases.
 - c. Kahawai are an Australasian species. In Australia, they range from Brisbane to Tasmania. They are also found at Lord Howe, Norfolk and Kermadec islands. In New Zealand they range from the Three Kings Islands to Otago and northern Fiordland, and the Chatham Islands. The known depth range is 0–150 m, though surface schools often occur over deeper water.

- d. Summer, for the purposes of NABIS, is defined as being the months of January, February and March. This definition is based on research regarding the spatial and temporal variability of sea surface temperature in the New Zealand region (Uddstrom and Oien 1999).

5. References

The following sources provided useful information on the distribution of this species. This is not an exhaustive list of all references to the species.

- Anderson, O.F.; Bagley, N.W.; Hurst, R.J.; Francis, M.P.; Clark, M.R.; McMillan, P.J. (1998). Atlas of New Zealand fish and squid distributions from research bottom trawls. *NIWA Technical Report 42*. 303 p.
- Francis, M.P. (1991). Additions to the fish faunas of Lord Howe, Norfolk and Kermadec Islands, Southwest Pacific Ocean. *Pacific Science 45*: 204-220.
- Francis, M.P. (1993). Checklist of the coastal fishes of Lord Howe, Norfolk, and Kermadec Islands, Southwest Pacific Ocean. *Pacific Science 47*: 136-170.
- Francis, M.P. (1996). Geographic distribution of marine reef fishes in the New Zealand region. *New Zealand Journal of Marine and Freshwater Research 30*: 35-55.
- Graham, D.H. (1938). Fishes of Otago Harbour and adjacent seas with additions to previous records. *Transactions and Proceedings of the Royal Society of New Zealand 68*: 399-419.
- Graham, D.H. (1956). A treasury of New Zealand fishes. Second edition. Reed, Wellington. 424 p.
- Graham, J. (1963). The North Otago shelf fauna. Part III - Chordata, sub-phylum Gnathostomata. *Transactions of the Royal Society of New Zealand. Zoology 3*: 165-170.
- Hardy, G.S.; Grace, R.V.; Francis, M.P. (1987). Fishes observed at the Three Kings Islands, northern New Zealand. *Records of the Auckland Institute and Museum 24*: 243-250.
- Hurst, R.J.; Bagley, N.W.; Anderson, O.F.; Francis, M.P.; Griggs, L.H.; Clark, M.R.; Paul, L.J.; Taylor, P.R. (2000a). Atlas of juvenile and adult fish and squid distributions from bottom and midwater trawls and tuna longlines in New Zealand waters. *NIWA Technical Report 84*. 162 p.
- Hurst, R.J.; Stevenson, M.L.; Bagley, N.W.; Griggs, L.H.; Morrison, M.A.; Francis, M.P. (2000b). Areas of importance for spawning, pupping or egg-laying, and juveniles of New Zealand coastal fish. *Final Research Report for Ministry of Fisheries Research Project ENV1999/03, Objective 1*. 302 p.

- Paulin, C. (1993). Review of the Australasian fish family Arripididae (Percomorpha), with the description of a new species. *Australian Journal of Marine and Freshwater Research* 44: 459-471.
- Roberts, C.D. (1991). Fishes of the Chatham Islands, New Zealand: a trawl survey and summary of the ichthyofauna. *New Zealand Journal of Marine and Freshwater Research* 25: 1-19.
- Uddstrom, M.J.; Oien, N.A. (1999). On the use of high-resolution satellite data to describe the spatial and temporal variability of sea surface temperatures in the New Zealand region. *Journal of Geophysical Research. Oceans* 104 C9: 20729-20751.