



Lineage

Winter distribution of tarakihi

Winter, for the purposes of NABIS, is defined as being the months of July, August and September. This definition is based on research regarding the spatial and temporal variability of sea surface temperature in the New Zealand region (Uddstrom and Oien 1999).

1. Electronic databases were used to generate initial maps of species distribution.
 - a. Commercial fishing returns (larger vessels): **TCEPR** database. All records from 1 October 1989 to 30 September 2005 were extracted on 17 October 2005. Data were used to estimate mean annual catch and catch rate (kilograms per kilometre towed) in 0.25 degree rectangles. Only the top five species caught are reported on these forms so information on the absence of a species is not available.
 - b. Commercial fishing returns (smaller vessels): **CELR** database. All records from 1 October 1989 to 30 June 2003 were extracted on 15–17 July 2003. Data were used to estimate mean annual catch in statistical areas. Information from statistical areas 1–10 was down-weighted because of likely mis-recording of Fishstock instead of statistical area. Only the top five species caught are reported on these forms so information on the absence of a species is not available.
 - c. Scientific observer records from larger vessels: **obs** database. All records from 1 March 1990 to 30 September 2005 and stored in the new data format were extracted on 20 October 2005. Data were used to estimate mean annual catch and catch rate (kilograms per kilometre towed), and proportion of tows that caught the species, in 0.25 degree rectangles.
 - d. Research bottom trawl records: **fish_comm** database. This database is a groomed version of the research trawl database **trawl**. All records from 2 September 1978 to 30 September 2005 were extracted on 19 May 2006. Data were used to estimate total catch, proportion of tows that caught the species, and catch rate (kilograms per kilometre towed) in 0.25 degree rectangles.
 - e. Russian research bottom trawl records: **trawl** database. These data are a subset of the research trawl database **trawl**. All records were extracted on 9 August 2003. Data were used to determine the presence of this species north of 37 °S. Because the data are old (the most recent record was 1987), and there are problems with species identifications, these data were given low weighting.
 - f. Recreational fishing database: **rec_data**. All records were extracted on 24 July 2003. Data were used to determine the presence of a species in a variety of statistical reporting areas.

- g. Museum of New Zealand Te Papa records of this species based on voucher specimens held in their collection were searched for distributional information that added to the distributional ranges determined from other databases.
 - h. Databases of commercial tuna longline catches (**TLCER**), observer records from tuna longlines (**I_line**), and aerial sightings (**aer_sight**) were not used as they contained no records of this species, or the number of records was too small to provide useful additional distributional information.
- 2. Literature sources were searched for distributional information that added to the distributional ranges determined from databases.
 - a. Unpublished electronic bibliography of New Zealand fishes compiled by L. J. Paul and held on a NIWA computer.
 - b. Aquatic Sciences and Fisheries Abstracts.
 - c. *New Zealand Professional Fisherman* and *Seafood New Zealand* for 1986–2002.
 - d. *New Zealand Fishing News* for 1998–2002.
 - e. Scientific papers, unpublished reports and university theses available to the expert who prepared the distributional layers.
- 3. Other sources.
 - a. Nil.
- 4. Summary
 - a. Maps generated from the electronic databases were provided to an expert scientist who integrated this information with other information from the literature, and expert opinion, and produced hand-drawn distributional zones on a template map containing depth contours at 250 m, 500 m, and 1000 m. These maps were then digitised and imported into a GIS software package as layers. The areas of the zones were calculated, and the layers were linked to attribute and metadata files.
 - b. The primary sources of distributional data for tarakihi were TCEPR, CELR, and fish_comm databases.
 - c. Tarakihi is an Indo-Pacific species, occurring in the south-western Atlantic Ocean, at St. Paul and Amsterdam islands in the Indian Ocean, southern Australia, including Tasmania, and in New Zealand. In New Zealand, tarakihi range from the Three Kings Islands to the southern edge of the Snares Shelf, and the Chatham islands. Hot spots for tarakihi in winter occur east of North Cape, north Taranaki, East Cape and Bay of Plenty, and the Chatham Islands. Their depth range is 10–500 m.
 - d. A closely-related, undescribed species (*Nemadactylus* sp., king tarakihi) occurs in northern New Zealand, and has been confused with tarakihi in the past (Smith et al. 1996). Records from the Wanganella Bank and Norfolk Island (Clark 1988, Francis 1993) may be the undescribed species and have been ignored.

5. References

The following sources provided useful information on the distribution of this species. This is not an exhaustive list of all references to the species.

Anderson, O.F.; Bagley, N.W.; Hurst, R.J.; Francis, M.P.; Clark, M.R.; McMillan, P.J. (1998). Atlas of New Zealand fish and squid distributions from research bottom trawls. *NIWA Technical Report 42*. 303 p.

Beentjes, M.P.; Bull, B.; Hurst, R.J.; Bagley, N.W. (2002). Demersal fish assemblages along the continental shelf and upper slope of the east coast of the South Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 36: 197-223.

Clark, M.R. (1988). Records of fishes from the West Norfolk Ridge, north-west of New Zealand. *New Zealand Journal of Zoology* 15: 415-421.

Francis, M.P. (1993). Checklist of the coastal fishes of Lord Howe, Norfolk, and Kermadec Islands, Southwest Pacific Ocean. *Pacific Science* 47: 136-170.

Hardy, G.S. (1986). An annotated list of fishes from the Snares Islands, New Zealand. *Mauri Ora* 13: 23-34.

Hardy, G.S.; Grace, R.V.; Francis, M.P. (1987). Fishes observed at the Three Kings Islands, northern New Zealand. *Records of the Auckland Institute and Museum* 24: 243-250.

Hurst, R.J.; Bagley, N.W.; Anderson, O.F.; Francis, M.P.; Griggs, L.H.; Clark, M.R.; Paul, L.J.; Taylor, P.R. (2000a). Atlas of juvenile and adult fish and squid distributions from bottom and midwater trawls and tuna longlines in New Zealand waters. *NIWA Technical Report 84*. 162 p.

Hurst, R.J.; Stevenson, M.L.; Bagley, N.W.; Griggs, L.H.; Morrison, M.A.; Francis, M.P. (2000b). Areas of importance for spawning, pupping or egg-laying, and juveniles of New Zealand coastal fish. *Final Research Report for Ministry of Fisheries Research Project ENV1999/03, Objective 1*. 302 p.

Kendrick, T.H.; Francis, M.P. (2002). Fish assemblages in the Hauraki Gulf, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 36: 699-717.

Roberts, C.D. (1991). Fishes of the Chatham Islands, New Zealand: a trawl survey and summary of the ichthyofauna. *New Zealand Journal of Marine and Freshwater Research* 25: 1-19.

Smith, P.J.; Roberts, C.D.; McVeagh, S.M.; Benson, P.G. (1996). Genetic evidence for two species of tarakihi (Teleostei: Cheilodactylidae: *Nemadactylus*) in New Zealand waters. *New Zealand Journal of Marine and Freshwater Research* 30: 209-220.

Uddstrom, M.J.; Oien, N.A. (1999). On the use of high-resolution satellite data to describe the spatial and temporal variability of sea surface temperatures in

the New Zealand region. *Journal of Geophysical Research. Oceans* 104 C9:
20729-20751.